

# Letzo 2.5 mg Film-Coated Tablets

Letrozole

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

## What is in this leaflet

- 1 What Letzo is and what it is used for
- 2 What you need to know before you take Letzo
- 3 How to take Letzo
- 4 Possible side effects
- 5 How to store Letzo
- 6 Contents of the pack and other information

**ROWEX®**

## 1 What Letzo is and what it is used for

### What Letzo is and how it works

Letzo contains an active substance called letrozole. It belongs to a group of medicines called aromatase inhibitors. It is a hormonal (or “endocrine”) breast cancer treatment. Growth of breast cancer is frequently stimulated by oestrogens which are female sex hormones. Letzo reduces the amount of oestrogen by blocking an enzyme (“aromatase”) involved in the production of oestrogens and therefore may block the growth of breast cancer that needs oestrogens to grow. As a consequence tumour cells slow or stop growing and/or spreading to other parts of the body.

### What Letzo is used for

Letzo is used to treat breast cancer in women who have gone through menopause i.e. cessation of periods.

It is used to prevent cancer from happening again. It can be used as first treatment before breast cancer surgery in case immediate surgery is not suitable or it can be used as first treatment after breast cancer surgery or following five years treatment with tamoxifen. Letzo is also used to prevent breast tumour spreading to other parts of the body in patients with advanced breast cancer.

If you have any questions about how Letzo works or why this medicine has been prescribed for you, ask your doctor.

## 2 What you need to know before you take Letzo

Follow all the doctor’s instructions carefully. They may differ from the general information in this leaflet.

### Do not take Letzo

- if you are allergic to letrozole or to any of the other ingredients of this medicine (listed in section 6)
- if you still have periods, i.e. if you have not yet gone through the menopause
- if you are pregnant
- if you are breast-feeding.

If any of these conditions apply to you, **do not take this medicine and talk to your doctor.**

### Warnings and precautions

Talk to your doctor or pharmacist before taking Letzo

- if you have a severe kidney disease
- if you have a severe liver disease
- if you have a history of osteoporosis or bone fractures (see also “Follow-up during Letzo treatment” in section 3).

If any of these conditions apply to you, tell your doctor. Your doctor will take this into account during your treatment with Letzo.

### Children and adolescents

Children and adolescents should not use this medicine.

### Older people (age 65 years and over)

People aged 65 years and over can use this medicine at the same dose as for other adults.

### Other medicines and Letzo

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

### Pregnancy, breast-feeding and fertility

- You should only take Letzo when you have gone through the menopause.  
However, your doctor should discuss with you about using effective contraceptive, as you may still have the potential to become pregnant during treatment with Letzo.
- If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, you must not take Letzo as it may harm your baby. Ask your doctor or pharmacist for advice before taking this medicine.

### Driving and using machines

If you feel dizzy, tired, drowsy or generally unwell, do not drive or operate any tools or machines until you feel normal again.

### Letzo contains lactose

Letzo contains lactose (milk sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## 3 How to take Letzo

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose is one tablet of Letzo to be taken once a day. Taking Letzo at the same time each day will help you remember when to take your tablet. The tablet can be taken with or without food and should be swallowed whole with a glass of water or another liquid.

### How long to take Letzo

Continue taking Letzo every day for as long as your doctor tells you. You may need to take it for months or even years. If you have any questions about how long to keep taking Letzo, talk to your doctor.

### Follow-up during Letzo treatment

You should only take this medicine under strict medical supervision. Your doctor will regularly monitor your condition to check whether the treatment is having the right effect. Letzo may cause thinning or wasting of your bones (osteoporosis) due to the reduction of oestrogens in your body. Your doctor may decide to measure your bone density (a way of monitoring for osteoporosis) before, during and after treatment.

### If you take more Letzo than you should

If you have taken too much Letzo, or if someone else accidentally takes your tablets, contact a doctor or hospital for advice immediately. Show them the pack of tablets. Medical treatment may be necessary.

### If you forget to take Letzo

- If it is almost time for your next dose (e.g. within 2 or 3 hours), skip the dose you missed and take your next dose when you are meant to.
- Otherwise, take the dose as soon as you remember, and then take the next tablet as you would normally.

- Do not take a double dose to make up for the one that you missed.

### If you stop taking Letzo

Do not stop taking Letzo unless your doctor tells you to. See also the section above "How long to take Letzo".

## 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Most of the side effects are mild to moderate and will generally disappear after a few days to a few weeks of treatment.

Some of these side effects, such as hot flushes, hair loss or vaginal bleeding, may be due to the lack of oestrogens in your body.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

### Some side effects could be serious:

**Rare or uncommon side effects** (may affect up to 1 in 100 people):

- Weakness, paralysis or loss of feeling in any part of the body (particularly arm or leg), loss of coordination, nausea, or difficulty speaking or breathing (sign of a brain disorder, e.g. stroke).
- Sudden oppressive chest pain (sign of a heart disorder).
- Difficulty breathing, chest pain, fainting, rapid heart rate, bluish skin discoloration, or sudden arm, leg or foot pain (signs that a blood clot may have formed).
- Swelling and redness along a vein which is extremely tender and possibly painful when touched.
- Severe fever, chills or mouth ulcers due to infections (lack of white blood cells).
- Severe persistent blurred vision.

### If any of the above occurs, tell your doctor straight away.

You should also **inform the doctor straight away if you experience any of the following symptoms** during treatment with Letzo:

- Swelling mainly of the face and throat (signs of allergic reaction)
- Yellow skin and eyes, nausea, loss of appetite, dark-coloured urine (signs of hepatitis)
- Rash, red skin, blistering of the lips, eyes or mouth, skin peeling, fever (signs of skin disorder).

**Some side effects are very common:** may affect more than 1 in 10 people.

- Hot flushes
- Increased level of cholesterol (hypercholesterolaemia)
- Fatigue
- Increased sweating
- Pain in bones and joints (arthralgia).

If any of these affects you severely, tell your doctor.

**Some side effects are common:** may affect up to 1 in 10 people.

- Skin rash
- Headache
- Dizziness
- Malaise (generally feeling unwell)
- Gastrointestinal disorders such as nausea, vomiting, indigestion, constipation, diarrhoea
- Increase in or loss of appetite
- Pain in muscles
- Thinning or wasting of your bones (osteoporosis), leading to bone fractures in some cases (see also "Follow-up during Letzo treatment" in section 3)
- Swelling of arms, hands, feet, ankles (oedema)
- Depression
- Weight increase
- Hair loss
- Raised blood pressure (hypertension)
- Abdominal pain
- Dry skin
- Vaginal bleeding.

If any of these affects you severely, tell your doctor.

**Other side effects are uncommon:** may affect up to 1 in 100 people.

- Nervous disorders such as anxiety, nervousness, irritability, drowsiness, memory problems, somnolence, insomnia
- Impairment of sensation, especially that of touch
- Eye disorders such as blurred vision, eye irritation

- Palpitations, rapid heart rate
- Skin disorders such as itching (urticaria)
- Vaginal discharge or dryness
- Joint stiffness (arthritis)
- Breast pain
- Fever
- Thirst, taste disorder, dry mouth
- Dryness of mucous membranes
- Weight decrease
- Urinary tract infection, increased frequency of urination
- Cough
- Increased level of enzymes.

If any of these affects you severely, tell your doctor.

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Letzo

Keep this medicine out of the sight and reach of children.

Do not store above 30°C.

Store in the original package in order to protect from moisture.

Do not use this medicine after the expiry date which is stated on the blister and carton after EXP. The first two digits indicate the month and the last four digits indicate the year. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## 6 Contents of the pack and other information

### What Letzo contains

The active substance is letrozole. Each film-coated tablet contains 2.5 mg letrozole.

The other ingredients are lactose monohydrate, cellulose microcrystalline (E460), maize starch, sodium starch glycolate, magnesium stearate (E572) and silica colloidal anhydrous (E551). The coating is composed of hypromellose (E464), talc (E553b), macrogol 8000, titanium dioxide (E171) and iron oxide yellow (E172).

### What Letzo looks like and contents of the pack

Letzo is supplied as film-coated tablets. The film-coated tablets are dark-yellow and round. They are marked with "FV" on one side and "CG" on the other side.

Each blister pack contains 10, 14, 28, 30, 50, 56, 60, 84, 90, 98, 100, 100 (10 x 10), 112 or 120 tablets. Not all pack sizes may be marketed.

### Marketing Authorisation Holder and Manufacturer

**Marketing Authorisation Holder**  
Rowex Ltd., Bantry, Co. Cork, Ireland.

**Manufacturer**  
Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, 39179 Barleben, Germany.

### This medicinal product is authorised in the Member States of the EEA under the following names:

The Netherlands	Letrozol Sandoz tablet 2,5 mg, filmomhulde tabletten
Germany	Letrozol HEXAL 2,5 mg Filmtabletten
Ireland	Letzo 2.5 mg Film-Coated Tablets

### This leaflet was last revised in 08/2016.

I.M. L/351a 08-16  
article no.