

Package leaflet: Information for the patient

Mirap DisTab 15 mg Orodispersible Tablets

Mirtazapine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Mirap DisTab is and what it is used for
2. What you need to know before you take Mirap DisTab
3. How to take Mirap DisTab
4. Possible side effects
5. How to store Mirap DisTab
6. Contents of the pack and other information

1. What Mirap DisTab is and what it is used for

Mirap DisTab is one of a group of medicines called **antidepressants**.

Mirap DisTab is used to treat depressive illness in adults.

Mirap DisTab will take 1 to 2 weeks before it starts working. After 2 to 4 weeks you may start feeling better. You must talk to your doctor if you do not feel better or if you feel worse after 2 to 4 weeks. More information is in section 3 heading "When can you expect to start feeling better".

2. What you need to know before you take Mirap DisTab

Do not take Mirap DisTab:

- if you are **allergic** to mirtazapine or any of the other ingredients of this medicine (listed in section 6). If so, you must talk to your doctor as soon as you can before taking Mirap DisTab.
- if you are taking or have recently taken (within the last two weeks) medicines called monoamine oxidase inhibitors (MAO-Is).

Warnings and precautions

Talk to your doctor or pharmacist before taking Mirap DisTab.

Children and adolescents

Mirap DisTab should normally not be used for children and adolescents under 18 years because efficacy was not demonstrated. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Mirap DisTab for patients under 18 because he/she decides that this is in their best interests. If your doctor has prescribed Mirap DisTab for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any of the symptoms listed above develop or worsen when patients under 18 are taking Mirap DisTab. Also, the long-term safety effects concerning growth, maturation and cognitive and behavioural development of Mirap DisTab in this age group have not yet been demonstrated. In addition, significant weight gain has been observed in this age category more often when treated with Mirap DisTab compared with adults.

Thoughts of suicide and worsening of your depression

If you are depressed you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
 - if you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant.
 - If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital straightaway.
- You may find it helpful to tell a relative or close friend** that you are depressed, and ask them to read this leaflet. You might ask them to tell you if they think your depression is getting worse, or if they are worried about changes in your behaviour.
- Also take special care with Mirap DisTab
- if you have, or have ever had one of the following conditions.
 - Tell your doctor about these conditions before taking Mirap DisTab, if not done previously.
 - **seizures** (epilepsy). If you develop seizures or your seizures become more frequent, stop taking Mirap DisTab and contact your doctor immediately;
 - **liver disease**, including jaundice. If jaundice occurs, stop taking Mirap DisTab and contact your doctor immediately;
 - **kidney disease**;
 - **heart disease**, or **low blood pressure**;
 - **schizophrenia**. If psychotic symptoms, such as paranoid thoughts become more frequent or severe, contact your doctor straightaway;
 - **manic depression** (alternating periods of feeling elated/overactivity and depressed mood). If you start feeling elated or over-excited, stop taking Mirap DisTab and contact your doctor immediately;
 - **diabetes** (you may need to adjust your dose of insulin or other antidiabetic medicines);
 - **eye disease**, such as increased pressure in the eye (glaucoma);
 - **difficulty in passing water** (urinating), which might be caused by an enlarged prostate.
 - **certain kinds of heart conditions** that may change your heart rhythm, a recent heart attack, heart failure, or take certain medicines that may affect the heart's rhythm.
 - if you develop signs of infection such as inexplicable high fever, sore throat and mouth ulcers.
 - Stop taking Mirap DisTab and consult your doctor immediately for a blood test.
- In rare cases these symptoms can be signs of disturbances in blood cell production in the bone marrow. While rare, these symptoms most commonly appear after 4-6 weeks of treatment.
- if you are an elderly person. You could be more sensitive to the side-effects of antidepressants.

Other medicines and Mirap DisTab

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Do not take Mirap DisTab in combination with:

- **monoamine oxidase inhibitors** (MAO inhibitors). Also, do not take Mirap DisTab during the two weeks after you have stopped taking MAO inhibitors. If you stop taking Mirap DisTab, do not take MAO inhibitors during the next two weeks either. Examples of MAO inhibitors are moclobemide, tranylcypromine (both are antidepressants) and selegiline (used for Parkinson's disease).

Take care when taking Mirap DisTab in combination with:

- **antidepressants such as SSRIs, venlafaxine and L-tryptophan, or triptans** (used to treat migraine), **tramadol** (a pain-killer), **linezolid** (an antibiotic), **lithium** (used to treat some psychiatric conditions) **methylen blue** (used to treat high levels of methemoglobin in the blood) **and St. John's Wort – Hypericum perforatum preparations** (a herbal remedy for depression). In very rare cases Mirap DisTab alone or the combination of Mirap DisTab with these medicines, can lead to a so-called serotonin syndrome. Some of the symptoms of this syndrome are: inexplicable fever, sweating, increased heart rate, diarrhoea, (uncontrollable) muscle contractions, shivering, overactive reflexes, restlessness, mood changes, and unconsciousness. If you get a combination of these symptoms, talk to your doctor immediately.
 - **the antidepressant nefazodone**. It can increase the amount of Mirap DisTab in your blood. Inform your doctor if you are using this medicine. It might be needed to lower the dose of Mirap DisTab, or when use of nefazodone is stopped, to increase the dose of Mirap DisTab again.
 - **medicines for anxiety or insomnia** such as benzodiazepines;
 - **medicines for schizophrenia** such as olanzapine;
 - **medicines for allergies** such as cetirizine;
 - **medicines for severe pain** such as morphine.
- In combination with these medicines Mirap DisTab can increase the drowsiness caused by these medicines.
- **medicines for infections**; medicines for bacterial infections (such as erythromycin); medicines for fungal infections (such as ketoconazole) and medicines for HIV/AIDS (such as HIV-protease inhibitors), **and drugs for stomach ulcers** (such as cimetide).
- In combination with Mirap DisTab these medicines can increase the amount of Mirap DisTab in your blood. Inform your doctor if you are using these medicines. It might be needed to lower the dose of Mirap DisTab, or when these medicines are stopped, to increase the dose of Mirap DisTab again.
- **medicines for epilepsy** such as carbamazepine and phenytoin;
 - **medicines for tuberculosis** such as rifampicin.
- In combination with Mirap DisTab these medicines can reduce the amount of Mirap DisTab in your blood. Inform your doctor if you are using these medicines. It might be needed to increase the dose of Mirap DisTab, or when these medicines are stopped to lower the dose of Mirap DisTab again.
- **medicines to prevent blood clotting** such as warfarin.
- Mirap DisTab can increase the effects of warfarin on the blood. Inform your doctor if you are using this medicine. In case of combination it is advised that a doctor monitors your blood carefully.
- **medicines that may affect the heart's rhythm** such as certain antibiotics and some anti-psychotics.

Mirap DisTab with food and alcohol:

You may get drowsy if you drink alcohol while you are taking Mirap DisTab.

You are advised not to drink any alcohol.

You can take Mirap DisTab with or without food.

Pregnancy and breast-feeding:

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Limited experience with Mirap DisTab administration to pregnant women does not indicate an increased risk. However, caution should be exercised when used during pregnancy.

If you use Mirap DisTab until, or shortly before birth, your baby should be supervised for possible adverse effects. When taken during pregnancy, similar drugs (SSRIs) may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Driving and using machines:

Mirap DisTab can affect your concentration or alertness. Make sure these abilities are not affected before you drive or operate machinery. If your doctor has prescribed Mirap DisTab for a patient under 18 years make sure the concentration and alertness is not affected before participation in traffic (e.g. on bicycle).

Mirap DisTab contains aspartame:

Mirap DisTab orodispersible tablets contain aspartame, a source of phenylalanine. May be harmful for people with phenylketonuria.

Mirap DisTab contains sulphites:

This medicine contains a very low amount of sulphites. This may rarely cause severe allergic (hypersensitivity) reactions and bronchospasm.

3. How to take Mirap DisTab

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to take

The recommended starting dose is 15 or 30 mg every day. Your doctor may advise you to increase your dose after a few days to the amount that is best for you (between 15 and 45 mg per day). The dose is usually the same for all ages. However, if you are an elderly person or if you have renal or liver disease, your doctor may adapt the dose.

When to take Mirap DisTab

→ Take Mirap DisTab as the same time each day.

It is best to take Mirap DisTab as a single dose before you go to bed. However your doctor may suggest to split your dose of Mirap DisTab – once in the morning and once at night-time before you go to bed. The higher dose should be taken before you go to bed.

Take the orodispersible tablet as follows

Take your tablets orally.

1. Do not crush the orodispersible tablet

In order to prevent crushing the orodispersible tablet, do not push against the tablet pocket (Figure A).



Fig. A.

2. Tear off one tablet pocket

Each blister contains tablet pockets, which are separated by perforations. Tear off one tablet pocket along the dotted lines (Figure 1).




Fig. 1.

3. Peel off the lid

Carefully peel off the lidding foil, starting in the corner indicated by the arrow (Figures 2 and 3).



Fig. 2.




Fig. 3.

4. Take out the orodispersible tablet

Take out the orodispersible tablet with dry hands and place it on the tongue (Figure 4).




Fig. 4

It will rapidly disintegrate and can be swallowed without water.

When can you expect to start feeling better

Usually Mirap DisTab will start working after 1 to 2 weeks and after 2 to 4 weeks you may start to feel better. It is important that, during the first few weeks of the treatment, you talk with your doctor about the effects of Mirap DisTab:

→ 2 to 4 weeks after you have started taking Mirap DisTab, talk to your doctor about how this medicine has affected you.

If you still don't feel better, your doctor may prescribe a higher dose. In that case, talk to your doctor again after another 2 to 4 weeks. Usually you will need to take Mirap DisTab until your symptoms of depression have disappeared for 4 to 6 months.

If you take more Mirap DisTab than you should

→ If you or someone else has taken too much Mirap DisTab, call a doctor straightaway. The most likely signs of an overdose of Mirap DisTab (without other medicines or alcohol) are **drowsiness, disorientation and increased heart rate**. The symptoms of a possible overdose may include changes to your heart rhythm (fast, irregular heartbeat) and/or fainting which could be symptoms of a life-threatening condition known as Torsade de Pointes.

If you still don't feel better, your doctor may prescribe a higher dose. In that case, talk to your doctor again after another 2 to 4 weeks. Usually you will need to take Mirap DisTab until your symptoms of depression have disappeared for 4 to 6 months.

If you forget to take Mirap DisTab

If you are supposed to take your dose **once a day**

- Do not take a double dose to make up for a forgotten dose. Take your next dose at the normal time.

If you are supposed to take your dose **twice a day**

- if you have forgotten to take your morning dose, simply take it together with your evening dose.
- if you have forgotten to take your evening dose, do not take it with the next morning dose; just skip it and continue with your normal morning and evening doses.
- if you have forgotten to take both doses, do not attempt to make up for the missed doses. Skip both doses and continue the next day with your normal morning and evening doses.

If you stop taking Mirap DisTab

→ Only stop taking Mirap DisTab in consultation with your doctor.

If you stop too early, your depression might come back. Once you are feeling better, talk to your doctor. Your doctor will decide when treatment can be stopped.

Do not suddenly stop taking Mirap DisTab, even when your depression has lifted. If you suddenly stop taking Mirap DisTab you may feel sick, dizzy, agitated or anxious, and have headaches. These symptoms can be avoided by stopping gradually. Your doctor will tell you how to decrease the dose gradually.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience any of the following serious side effects, stop taking mirtazapine and tell your doctor immediately.

Uncommon (may affect up to 1 in 100 people):

- feeling elated or emotionally 'high' (mania)

Rare (may affect up to 1 in 1,000 people):

- yellow colouring of eyes or skin; this may suggest disturbance in liver function (jaundice)

Not known (frequency cannot be estimated from the available data):

- signs of infection such as sudden unexplainable high fever, sore throat and mouth ulcers (agranulocytosis). In rare cases mirtazapine can cause disturbances in the production of blood cells (bone marrow depression). Some people become less resistant to infection because mirtazapine can cause a temporary shortage of white blood cells (granulocytopenia). In rare cases mirtazapine can also cause a shortage of red and white blood cells, as well as blood platelets (aplastic anemia), a shortage of blood platelets (thrombocytopenia) or an increase in the number of white blood cells (eosinophilia).
- epileptic attack (convulsions)
- a combination of symptoms such as inexplicable fever, sweating, increased heart rate, diarrhoea, (uncontrollable) muscle contractions, shivering, overactive reflexes, restlessness, mood changes, unconsciousness and increased salivation. In very rare cases these can be signs of serotonin syndrome.
- thoughts of harming or killing yourself
- severe skin reactions (Stevens-Johnson Syndrome, toxic epidermal necrolysis)

Other possible side-effects with mirtazapine are:

Very common (may affect more than 1 in 10 people):

- increase in appetite and weight gain
- drowsiness or sleepiness
- headache
- dry mouth

Common (may affect up to 1 in 10 people):

- lethargy
- dizziness
- shakiness or tremor
- nausea
- diarrhoea
- vomiting
- constipation
- rash or skin eruptions (exanthema)
- pain in your joints (arthralgia) or muscles (myalgia)
- back pain
- feeling dizzy or faint when you stand up suddenly (orthostatic hypotension)
- swelling (typically in ankles or feet) caused by fluid retention (oedema)
- tiredness
- vivid dreams
- confusion
- feeling anxious
- sleeping problems

Uncommon (may affect up to 1 in 100 people):

- abnormal sensation in the skin e.g. burning, stinging, tickling or tingling (paraesthesia)
- restless legs
- fainting (syncope)
- sensations of numbness in the mouth (oral hypoaesthesia)
- low blood pressure
- nightmares
- feeling agitated
- hallucinations
- urge to move

Rare (may affect up to 1 in 1,000 people):

- muscle twitching or contractions (myoclonus)
- aggression
- abdominal pain and nausea; this may suggest inflammation of the pancreas (pancreatitis)

Not known (frequency cannot be estimated from the available data):

- abnormal sensations in the mouth (oral paraesthesia)
- swelling in the mouth (mouth oedema)
- swelling throughout the body (generalized oedema)
- localized swelling
- hyponatraemia
- inappropriate anti-diuretic hormone secretion
- severe skin reactions (dermatitis bullous, erythema multiforme)
- sleep walking (somnambulism)
- speech disorder

Other possible side effects with mirtazapine are

Increased creatine kinase blood levels, difficulty in passing urine and muscle pain, stiffness and/or weakness and darkening or discolouration of the urine.

Additional side effects in children and adolescents

In children under 18 years the following adverse events were observed commonly in clinical trials: significant weight gain, hives and increased blood triglycerides.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Mirap DisTab

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month.

Do not store above 30°C. Store in the original package in order to protect from moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Mirap DisTab contains

- The active substance is: Mirtazapine. Each orodispersible tablet contains 15 mg mirtazapine.
- The other ingredients are: mannitol (E421), povidone K30, crospovidone, silica colloidal anhydrous, aspartame (E951), calcium stearate, orange flavour [maltodextrin, natural and artificial flavourings, dl-alpha-tocopherol], peppermint flavour [maltodextrin, natural flavourings, dextrin, sulphites].

What Mirap DisTab looks like and contents of the pack

Orodispersible tablets: White to off-white, round, flat tablets with bevelled edges and plain on both sides.

Aluminium/Aluminium blisters containing 6, 10, 18, 28, 30, 48, 84, 90 or 96 orodispersible tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Rowex Ltd., Bantry, Co. Cork, Ireland.

Manufacturers

Lek Pharmaceuticals d.d., Verovškova 57, 1526 Ljubljana, Slovenia.
Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, 39179 Barleben, Germany.
Salutas Pharma GmbH, Dieselstrasse 5, 70839 Gerlingen, Germany.
Sandoz GmbH, Biochemiestrasse 10, 6250 Kundl, Austria.

This medicinal product is authorised in the Member States of the EEA under the following names:

Austria:	Mirtazapin "Hexal" 15 mg - Schmelztabletten
Ireland:	Mirap DisTab 15 mg Orodispersible Tablets
Luxembourg:	Mirtazapin HEXAL 15 mg Schmelztabletten
The Netherlands:	Mirtazapine orodispergeerbare tablet 15 mg, orodispergeerbare tabletten 15 mg
Spain:	Mirtazapina Sandoz 15 mg comprimidos bucodispersables EFG

This leaflet was last revised in 02/2015.