

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Tradol 50 mg Effervescent Tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each effervescent tablet contains Tramadol Hydrochloride 50mg.

Excipient with known effect:

Each tablet contains 75mg lactose monohydrate and 231mg sodium

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Effervescent Tablet.

White to almost white, round biplanar effervescent tablets, flat, smell like orange, with a facet on both sides and a score notch on one side.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic Indications

Treatment of moderate to severe pain.

### 4.2 Posology and method of administration

The dose should be adjusted to the intensity of the pain and the sensitivity of the individual patient. The lowest effective dose for analgesia should generally be selected. The total daily dose of 400mg active substance should not be exceeded, except in special circumstances.

Unless otherwise prescribed, Tradol should be administered as follows:

Route of Administration: oral

*Adults and adolescents aged 12 years and over:*

Acute pain: an initial dose of 100mg is usually necessary. This can be followed by doses of 50 or 100mg at 4-6 hourly intervals and duration of treatment should be matched to clinical need.

Pain associated with chronic conditions: an initial dose of 50mg is advised and then titration according to pain severity. The need for continued treatment should be assessed at regular intervals as withdrawal symptoms and dependence have been reported (see section 4.4).

*Children*

Tramadol is not suitable for children below the age of 12 years.

*Geriatric patients:*

A dose adjustment is not usually necessary in patients up to 75 years without clinically manifest hepatic or renal insufficiency. In elderly patients over 75 years elimination may be prolonged. Therefore, if necessary, the dosage interval is to be extended according to the patient's requirements.

*Patients with renal Insufficiency/ dialysis and hepatic impairment:*

In patients with renal and/or hepatic insufficiency the elimination of tramadol is delayed. In these patients prolongation of the dosage intervals should be carefully considered according to the patient's requirements.

**Method of administration**

The tablets should be dissolved in about half a glass of water.

**Duration of administration**

Tramadol should under no circumstances be administered for longer than absolutely necessary. If long term pain treatment with Tramadol is necessary in view of the nature and severity of the illness, then careful and regular monitoring should be carried out (if necessary with breaks in the treatment) to establish whether and to what extent further treatment is necessary.

**4.3 Contraindications**

TRADOL is contraindicated

- in hypersensitivity to the active substance or to any of the excipients listed in section 6.1
- in acute intoxication with alcohol, hypnotics, analgesics, opioids or other psychotropic medicinal products
- in patients who are receiving monoamine oxidase inhibitors or who have taken them within the last 14 days (see section 4.5)
- in patients with epilepsy not adequately controlled by treatment.
- for use in narcotic withdrawal treatment.

**4.4 Special warnings and precautions for use**

TRADOL may only be used with particular caution in opioid-dependent patients, patients with head injury, shock, a reduced level of consciousness of uncertain origin, disorders of the respiratory centre or function, increased intracranial pressure, severe impairment of hepatic and renal function and in patients prone to convulsive disorders.

In patients sensitive to opiates the product should only be used with caution.

Convulsions have been reported in patients receiving Tramadol at the recommended dose levels. The risk may be increased when doses of Tramadol exceed the recommended upper daily limit (400mg). In addition tramadol may increase the seizure risk in patients taking other medicinal products that lowers the seizure threshold (see section 4.5). Patients with epilepsy or those susceptible to seizures should be only treated with tramadol if there are compelling circumstances.

Care should be taken when treating patients with respiratory depression, or if concomitant CNS depressant drugs are being administered (see section 4.5), or if the recommended dosage is significantly exceeded (see section 4.9) as the possibility of respiratory depression cannot be excluded in these situations. At therapeutic doses respiratory depression has infrequently been reported.

Tramadol has a low dependence potential. On long-term use tolerance, psychic and physical dependence may develop. In patients with a tendency to drug abuse or dependence, treatment with tramadol should only be carried out for short periods under strict medical supervision.

Tramadol is not suitable as a substitute in opioid-dependant patients. Although it is an opioid agonist, tramadol cannot suppress morphine withdrawal symptoms.

At therapeutic doses withdrawal symptoms have been reported at a frequency of 1 in 8,000. Reports of dependence and abuse have been less frequent. Because of this potential the clinical need for continued analgesic treatment should be reviewed regularly.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicinal product contains sodium and must be taken into consideration by patients on a controlled sodium diet.

#### 4.5 Interaction with other medicinal products and other forms of interaction

Tradol should not be combined with MAO inhibitors (see section 4.3).

In patients treated with MAO inhibitors in the last 14 days prior to the use of the opioid pethidine, life threatening interactions on the central nervous system, respiratory and cardiovascular functions have been observed. The same interactions with MAO inhibitors cannot be ruled out during treatment with tramadol.

Concomitant administration of TRADOL with other centrally depressant medicinal products, including alcohol may potentiate CNS effects (see section 4.8).

The results of pharmacokinetic studies have so far shown that on the concomitant or previous administration of cimetidine (enzyme inhibitor) clinically relevant interactions are unlikely to occur. Co-administration with cimetidine is associated with a small prolongation of the half-life of tramadol, but this is not clinically relevant. Simultaneous or previous administration of carbamazepine (enzyme inducer) may reduce the analgesic effect and shorten the duration of action as a result of a reduction in serum levels of tramadol and its active metabolite.

Other morphine derivatives (including anti-tussives, substitution treatments), benzodiazepines, barbiturates; increased risk of respiratory depression, that may be fatal in overdose.

Mixed agonists/antagonists (e.g. buprenorphine, nalbuphine, pentazocine); the analgesic effect of tramadol, which is a pure agonist, may be reduced and a withdrawal syndrome may occur.

Tramadol can induce convulsions and increase the potential for selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, antipsychotics and other seizure threshold-lowering medicinal products (such as bupropion, mirtazapine, tetrahydrocannabinol) to cause convulsions.

Concomitant therapeutic use of tramadol and serotonergic drugs, such as selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), MAO inhibitors (see section 4.3), tricyclic antidepressants and mirtazapine may cause serotonin toxicity. Serotonin syndrome is likely when one of the following is observed:

- Spontaneous clonus
- Inducible or ocular clonus with agitation or diaphoresis
- Tremor and hyperreflexia
- Hypertonia and body temperature >38°C and inducible or ocular clonus.

Withdrawal of the serotonergic drugs usually brings about a rapid improvement. Treatment depends on the type and severity of the symptoms.

Caution should be exercised when commencing and during concomitant treatment with tramadol and coumarin derivatives (e.g. warfarin) due to reports of increased INR, major bleeding and ecchymoses in some patients.

Other active substances known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation) probably also the metabolism of the active O-demethylated metabolite. The clinical importance of such an interaction has not been studied.

In a limited number of studies the pre-or postoperative application of the antiemetic 5-HT<sub>3</sub> antagonist ondansetron increased the requirements of tramadol in patients with post operative pain.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy

Animal studies with tramadol revealed at very high doses effects on organ development, ossification and neonatal mortality. Teratogenic effects were not observed. Tramadol crosses the placenta. There is inadequate evidence available on the safety of tramadol in human pregnancy. Therefore tramadol should not be used in pregnant women.

Tramadol-administered before or during birth-does not affect uterine contractility. In neonates it may induce changes in the respiratory rate which are usually not clinically relevant.

Chronic use during pregnancy may lead to neonatal withdrawal symptoms.

### Breast-feeding

During lactation about 0.1% of the maternal dose is secreted into the milk. Tramadol is not recommended during breast-feeding. After a single administration of tramadol it is not usually necessary to interrupt breast-feeding.

### Fertility

Post marketing surveillance does not suggest an effect of tramadol on fertility. Animal studies did not show an effect of tramadol on fertility.

## 4.7 Effects on ability to drive and use machines

Even when taken according to instructions, tramadol may cause effects such as somnolence and dizziness and therefore may impair the reactions of drivers and machine operators. This applies particularly in conjunction with other psychotropic substances, particularly alcohol.

## 4.8 Undesirable effects

The most commonly reported adverse reactions are nausea and dizziness, both occurring in more than 10% of patients.

The frequencies are defined as follows:

Very common  $\geq 1/10$ , common  $\geq 1/100$ ,  $< 1/10$ , uncommon  $\geq 1/1000$ ,  $< 1/100$ , rare  $\geq 1/10,000$ ,  $< 1/1000$ , very rare  $< 1/10,000$ , not known: cannot be estimated from the available data.

### **Cardiac disorders:**

Uncommon: cardiovascular regulation (palpitation, tachycardia). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.

Rare: bradycardia

### **Investigations:**

Rare: increase in blood pressure

### **Vascular disorders:**

Uncommon: cardiovascular regulation (postural hypotension or cardiovascular collapse). These adverse effects may occur especially on intravenous administration and in patients who are physically stressed.

### **Metabolism and nutrition disorders**

Rare: changes in appetite

Not known: hypoglycaemia

### **Respiratory, thoracic and mediastinal disorders:**

Rare: respiratory depression, dyspnoea

If the recommended doses are considerably exceeded and other centrally depressant substances are administered concomitantly (see Section 4.5), respiratory depression may occur.

Worsening of asthma has been reported, though a causal relationship has not been established.

#### **Nervous system disorders:**

Very common: dizziness

Common: headache, somnolence

Rare: speech disorders, paraesthesia, tremor, epileptiform convulsions, involuntary muscle contractions, abnormal coordination, syncope.

Convulsions occurred mainly after administration of high doses of tramadol or after concomitant treatment with medicinal products which can lower the seizure threshold (see Sections 4.4 and 4.5).

Not known: speech disorders

#### **Psychiatric disorders:**

Rare: hallucinations, confusion, sleep disturbance, delirium, anxiety and nightmares. Psychic adverse reactions may occur following administration of tramadol, which vary individually in intensity and nature (depending on personality and duration of treatment). These include changes in mood (usually elation, occasionally dysphoria), changes in activity (usually suppression, occasionally increase) and changes in cognitive and sensorial capacity (e.g. decision behaviour, perception disorders). Drug dependence may occur. Symptoms of withdrawal reactions, similar to those occurring during opiate withdrawal, may occur as follows: agitation, anxiety, nervousness, insomnia, hyperkinesia, tremor and gastrointestinal symptoms. Other symptoms that have very rarely been seen with tramadol discontinuation include: panic attacks, severe anxiety, hallucinations, paraesthesia, tinnitus and unusual CNS symptoms (i.e. confusion, delusions, depersonalisation, derealisation, paranoia).

#### **Eye disorders:**

Rare: miosis, mydriasis, blurred vision

#### **Gastrointestinal disorders:**

Very common: nausea

Common: constipation, dry mouth, vomiting

Uncommon: retching, gastrointestinal discomfort (a feeling of pressure in the stomach, bloating), diarrhoea.

#### **Skin and subcutaneous tissue disorders:**

Common: hyperhidrosis

Uncommon: dermal reactions (e.g. pruritus, rash, urticaria)

#### **Musculoskeletal and connective tissue disorders:**

Rare: motorial weakness.

#### **Hepatobiliary disorders:**

In a few isolated cases an increase in liver enzyme values have been reported in a temporal connection with the therapeutic use of tramadol

#### **Renal and urinary disorders:**

Rare: micturition disorders (dysuria and urinary retention)

#### **Immune system disorders**

Rare: Allergic reactions (e.g. dyspnoea, bronchospasm, wheezing, angioneurotic oedema) and anaphylaxis

#### **General disorders and administration site conditions**

Common: fatigue

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL-Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie).

## 4.9 Overdose

### Symptoms

In principle, on intoxication with tramadol symptoms similar to those of other centrally acting analgesics (opioids) are to be expected. These include in particular miosis, vomiting, cardiovascular collapse, consciousness disorders up to coma, convulsions and respiratory depression up to respiratory arrest.

### Treatment

The general emergency measures apply. Keep open the respiratory tract (aspiration!), maintain respiration and circulation depending on the symptoms. The antidote for respiratory depression is naloxone. In animal experiments naloxone had no effect on convulsions. In such cases diazepam should be given intravenously.

In case of intoxication orally, gastrointestinal decontamination with activated charcoal or by gastric lavage is only recommended within 2 hours after tramadol intake. Gastrointestinal decontamination at a later time point may be useful in case of intoxication with exceptionally large quantities or prolonged-release formulations.

Tramadol is minimally eliminated from the serum by haemodialysis or haemofiltration. Therefore treatment of acute intoxication with TRADOL with haemodialysis or haemofiltration alone is not suitable for detoxification.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

ATC code : Analgesics:

NO2AXO2: Other Opioids

TRADOL is a centrally acting opioid analgesic. It is a non selective pure agonist at mu, delta and kappa opioid receptors with a higher affinity for the mu receptor. Other mechanisms which may contribute to its analgesic effect are inhibition of neuronal reuptake of noradrenaline and enhancement of serotonin release.

Tramadol has an antitussive effect. In contrast to morphine, analgesic doses of tramadol over a wide range have no respiratory depressant effect. Also gastrointestinal motility is less affected. Effects on the cardiovascular system tend to be slight. The potency of tramadol is reported to be 1/10 (one tenth) to 1/6 (one sixth) that of morphine.

### 5.2 Pharmacokinetic properties

About 90% of tramadol is absorbed after oral administration.

The half-life of the terminal elimination phase ( $t_{1/2\beta}$ ) is  $6.0 \pm 1.5$  h in young volunteers.

Tramadol pharmacokinetics show little age-dependence, the minimal changes being therapeutically irrelevant. In patients above the age of 65 years the  $t_{1/2}$ ,  $\beta$  was  $6.5 \pm 1.7$  h on oral administration. In volunteers aged over 75 years,  $t_{1/2}$   $\beta$  was  $7.0 \pm 1.6$  h on oral administration. Since Tramadol is eliminated both metabolically and renally, the terminal half-life  $t_{1/2\beta}$  may be prolonged in impaired hepatic or renal function.

However, the increase in the  $t_{1/2\beta}$  value is relatively low if at least one of these organs is functioning normally. In patients with liver cirrhosis  $t_{1/2\beta}$ , Tramadol was a mean of  $13.3 \pm 4.9$  h, in patients with renal insufficiency (creatinine clearance  $< 5$  ml/min) it was  $11.0 \pm 3.2$  h.

### 5.3 Preclinical safety data

On repeated oral and parenteral administration of tramadol for 6 – 26 weeks in rats and dogs and oral administration for 12 months in dogs haematological, clinico-chemical and histological investigations showed no evidence of any substance-related changes. Central nervous manifestations only occurred after high doses considerably above the therapeutic range: restlessness, salivation, convulsions, and reduced weight gain. Rats and dogs tolerated oral doses of 20mg/kg and 10mg/kg body weight respectively, and dogs rectal doses of 20mg/kg body weight without any reactions.

In rats tramadol dosages from 50 mg/kg/day upwards caused toxic effects in dams and raised neonate mortality. In the offspring retardation occurred in the form of ossification disorders and delayed vaginal and eye opening. Male fertility was not affected. After higher doses (from 50 mg/kg/day upwards) females exhibited a reduced pregnancy rate. In rabbits there were toxic effects in dams from 125 mg/kg upwards and skeletal anomalies in the offspring. In some *in vitro* test systems there was evidence of mutagenic effects. *In vivo* studies showed no such effects. According to knowledge gained so far, tramadol can be classified as non-mutagenic.

Studies on the tumourigenic potential of tramadol hydrochloride have been carried out in rats and mice. The study in rats showed no evidence of any substance-related increase in the incidence of tumours. In the study in mice there was an increased incidence of liver cell adenomas in male animals (a dose-dependent, non-significant increase from 15 mg/kg upwards) and an increase in pulmonary tumours in females of all dosage groups (significant, but not dose-dependent).

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Anhydrous Citric Acid  
Sodium Hydrogen Carbonate  
Anhydrous Sodium Sulphate  
Lactose Monohydrate  
Macrogol 6000  
Anhydrous Sodium Carbonate  
Povidone  
Sodium Cyclamate  
Orange Flavour 1013042  
Simethicone Emulsion  
Magnesium stearate

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

4 years.

### 6.4 Special precautions for storage

Do not store above 25°C.  
Keep the container tightly closed in order to protect from moisture.

### 6.5 Nature and contents of container

TRADOL Effervescent Tablets are packed into tablet tubes made of white polypropylene. The tubes which contain either 10 or 15 tablets are closed with air-sac stoppers made of white polyethylene, fitted with Silica gel as siccative and a cover disc of carton.  
TRADOL Effervescent tablets 50 mg will be available in pack sizes of 5 x 10 tablets and 2 x 15 tablets.  
Not all pack sizes may be marketed.

### 6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

**7 MARKETING AUTHORISATION HOLDER**

Rowex Ltd.  
Bantry  
Co. Cork

**8 MARKETING AUTHORISATION NUMBER**

PA0711/029/002

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 25 June 1999

Date of last renewal: 25 June 2009

**10 DATE OF REVISION OF THE TEXT**

November 2015